

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Type 2 diabetes mellitus is associated with disruptions as a function of metabolic control.

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/76024> since 2017-06-20T16:07:26Z

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Type 2 diabetes mellitus is associated with disruptions as a function of metabolic control

M. Trento¹, F. Riganti¹, M. Tomelini¹, P. Passera, F. Broglio¹, E. van Cauter², M. Porta¹, E. Ghigo¹

¹Department of Internal Medicine, University of Turin, Italy; ²Department of Medicine, University of Chicago, IL, USA

Background and aims Sleep disturbances may be an independent risk factor for glucose intolerance and, in type 2 diabetes mellitus (T2DM), sleep disruption was shown to correlate with metabolic control. We aimed at evaluating sleep quality in non-obese T2DM patients treated with oral agents and without complications. **Methods** Sleep recording was performed in 25 middle-aged T2DM patients and 18 age- and sex-matched healthy controls subjects (CS) by wrist actigraphy (Actiwatch, MiniMitterCo, USA) on three consecutive working days in free living conditions. Data were analysed using ActiwareSleepSoftware. **Results** There were no significant differences between T2DM and CS in terms of time in bed (mean±SD 7h40'±0h53' vs 7h36'±1h06'), actual sleep time (6h25'±0h52' vs 6h36'±1h03'), sleep efficiency (83.77±6.92 vs 86.81±4.19%) and sleep latency (11'±11' vs 09'±07'). However, T2DM patients showed lower (p<0.05) actual sleep percentage (88.77±4.56 vs 91.27±3.15%), higher (p<0.01) fragmentation index (21.10±8.14 vs 15.26±3.90), higher (p<0.05) total activity score during night-time (7822±3596 vs 6339±3672) and increased (p<0.01) moving time percentage (11.26±5.09 vs 7.88±1.50%). In T2DM, HbA1c correlated negatively with sleep efficiency (r=-0.624; p<0.01) and actual sleep percentage (r=-0.585; p<0.01) and positively with fragmentation index (r=0.576; p<0.01), moving time percentage (r=0.651; p<0.01) and mean activity score during night-time (r=0.493; p<0.05). **Discussion** These data suggest that T2DM is associated with quantitative and qualitative sleep disruption which correlate with metabolic control, even in the absence of complications.